



Lyme Disease

Categories : [Health Issues](#)

Lyme disease is a tick-borne bacterial infection transmitted to humans by the bite of infected Ixodes Scapularis ticks, also known as black-legged or deer ticks, and are typically found in grassy and overgrown areas. Ticks do not jump, fly or drop out of trees, and usually are picked up on the lower legs and crawl up the body to find a feeding spot. The bite is usually painless, so people do not know they have been bitten. Most people become infected during the months of May-June when ticks are most active.

To prevent exposure to tick bites, use the following protective measures, and find additional tips and resources at the Connecticut Department of Public Health's: "Lyme Disease" website page, <https://portal.ct.gov/en/DPH/Infectious-Diseases/Tickborne/Lyme-Disease>:

- Avoid tall grass and over-grown areas
- When hiking stay in the middle of trails
- Consider using tick repellent
- Tuck pant leg into socks
- Wear long-sleeved shirts and closed shoes
- Wear light-colored clothing to see the ticks easier for removal
- Use insect repellent to help reduce the chance of tick bites (DEET is the primary active ingredient to repel ticks, and must be used as directed on the container)
- Examine yourself, your children, and pets for ticks when returning indoors
- When returning home after an outing, shower using a washcloth or puff to remove unattached ticks
- Dry your clothes before washing if you've been outside, take off clothes and throw them in the dryer on high heat for at least 20 minutes
- Talk to your veterinarian to find out how to protect your pets from tick bites, and the roll of vaccine for dogs

****Since Lyme disease is a multi-systemic illness there are a multitude of Lyme disease symptoms including****

Early signs and symptoms – generally appear 2-32 days after bite by an infected deer tick:

- Flu-like symptoms, including muscle aches, fatigue, and fever
- Expanding red rash (Note: many people do not get the red bull's eye expanding rash)

Other symptoms include:

- Other rashes on parts of body
- Facial or Bell's Palsy
- Heart palpitations
- Dizziness
- Joint pain & swelling

NOTE: There are a variety of transmittable diseases caused by ticks, besides Lyme disease, that have also been reported in Connecticut. For more information on these other disease names and symptoms, visit the Connecticut Department of Public Health's website page at: <http://www.portal.ct.gov/DPH/Infectious-Diseases/Tickborne/Other-Tick-borne-Diseases>



For additional information and resources on this disease, see the following:

- Centers for Disease Control: <http://www.cdc.gov/lyme/>
- Lymedisease.org: <https://www.lymedisease.org/news-blogs/news/>
- Lyme Disease Association: <https://www.lymediseaseassociation.org/>

TICK IDENTIFICATION AND TESTING IN CONNECTICUT

Ticks can be taken to your municipal health department or regional health district for identification and testing. The health department will deliver ticks to the Connecticut Agricultural Experiment Station, <http://www.ct.gov/caes/cwp/view.asp?a=2837&q=378220> for identification and testing. Individuals can also take ticks to the Connecticut Agricultural Experiment Station in New Haven for identification only.

TO FIND SUPPORT GROUPS FOR PEOPLE WITH LYME DISEASE IN CONNECTICUT'S COMMUNITY SERVICES DATABASE

Search by service term: [Health/Disability Related Support Groups*Lyme Disease](#)

-

SOURCES: Lyme Disease Association; Connecticut Department of Public Health; United States Centers for Disease Control

PREPARED BY: 211/rj

CONTENT LAST REVIEWED: April2019